

**ANTI-HYPERGLYCEMIC EFFECT OF SERPENTINA  
(*Andrographis paniculata*) LEAF EXTRACT IN  
ALLOXAN-INDUCED DIABETIC MICE**

**Rita B. Daliwag**  
**Michelle Anne Bunagan**  
**Mary Grace Ventura**  
*St. Paul University Philippines*  
*rdaliwag@spup.edu.ph*

**ABSTRACT**

*The study evaluated the anti-hyperglycemic effect of Serpentina leaf extract in alloxan-induced diabetic mice. Specifically, it aimed to determine whether there is a significant difference in the blood sugar level of diabetic mice when subjected to the test drug and the control. Alloxan monohydrate was administered to the test animals with single intraperitoneal injection at a dose rate of 125mg/kg once a day for six days, and on the seventh day, their blood sugar level was evaluated for individual glucose levels. Serpentina leaf extract was administered at doses of 14.1 ml/kg (Treatment 1), 28.2 ml/kg (Treatment 2) and 47 ml/kg (Treatment 3). Metformin was used as the positive control. The effect was monitored after 3 and 24 hours after oral medication with the test drug and the control. Results showed a mean difference of reduction levels after 3 hours at -44.7, 240.25, 140.5 and 18.75, and at 24 hours at 193.75, 233.75, 264.5 and 93 in the blood sugar levels of test animals treated with Serpentina and the control drug. The study further reveals that a dose of 28.2 ml/kg Serpentina leaf extract showed significant reduction of blood sugar levels in hyperglycemic mice sustained after three (3) hours and twenty-four (24) hours post treatment at 0.05 level of significance.*

**Keywords:** *anti-hyperglycemic, alloxan monohydrate, Serpentina, intraperitoneal*

## INTRODUCTION

Hyperglycemia is a widespread disorder, which is characterized by elevated plasma glucose concentrations resulting from insufficient insulin and insulin resistance. This could lead to the disease called diabetes mellitus.

Diabetes mellitus is a serious health problem with continuously increasing number of incidence and mortality. If not cured, this will eventually lead to chronic complications causing other diseases, infections, and serious organ damage.

Plants represent the richest source of inspiration for the identification of novel scaffold structures that can serve as the basis for drug design. Plants with antidiabetic properties, in particular are generating a lot of interest primarily because the number of diabetics worldwide are fast reaching epidemic proportions, with the total number of people with diabetes projected to rise from 171 million in 2000 to 366 million in 2030.

*Andrographis paniculata* is a plant that has been effectively used in traditional Asian medicines for three centuries. It is perceived “blood purifying” property results in its use in diseases where blood abnormalities are considered causes of disease, such as skin eruptions, boils, scabies, and chronic undetermined fevers (Kumar et al., 2012).

The present study evaluates the anti-hyperglycemic effect of *Andrographis paniculata* (Serpentina) leaf extract to alloxan-induced diabetic mice. This is to confirm the folkloric claim that the leaves of Serpentina can lower blood glucose levels.

### Objectives of the Study

This study aimed to determine the efficacy of the leaf extract of Serpentina (*Andrographis paniculata*) as anti-hyperglycemic agent. Specifically, it sought to answer the following questions:

1. What are the blood sugar levels of the mice after induction with

alloxan?

2. Is there a significant difference in the blood sugar level of the mice before and after treatment with metformin?
3. Is there a significant difference in the blood sugar level of the mice before and after the administration of the *Serpentina* leaf extract?
4. Is there a significant difference in the anti-hyperglycemic effect of *Serpentina* leaf extract and metformin?
5. What exact volume of the *Serpentina* leaf extract is effective as anti-hyperglycemic?

### **Hypotheses**

1. There is no significant difference in the blood sugar level of the mice before and after treatment with alloxan.
2. There is no significant difference in the blood sugar levels of the mice before and after the administration of metformin.
3. There is no significant difference in the blood sugar level of the mice before and after the administration of the leaf extract.
4. There is no significant difference in the anti-hyperglycemic effect of *Serpentina* and metformin.

## **METHODOLOGY**

### **Collection of Plant Material**

Leaves of *Serpentina* were collected from Barangay Cataggaman Nuevo in Tuguegarao City, Cagayan, Philippines. A sample was brought to the National Museum in Manila for authentication.

### **Preparation of Plant Extract**

Leaves of *Serpentina* were weighed, washed, and dried. Fifty grams (50g) of the leaves were placed in 500ml of 40% ethyl alcohol and allowed to stand for 3 days with frequent agitation, until soluble matter was dissolved. The mixture was strained, the marc (damp solid material) was pressed, and the combined liquids clarified by filtration after standing. The filtered extracts were processed by a rotary evaporator for the removal of the excess solvent in 70°C. The extract

was kept in a clean and sterilized container and refrigerated till further use.

### **Preparation of Diabetic Mice**

The test animals were acclimatized for one week prior to the administration of the different doses of Alloxan monohydrate. Alloxan monohydrates were administered to the 16 normal mice at 125 mg/kg once a day for six days. On the seventh day, the blood glucose level obtained by the tail cut technique was read with accu-check glucometer. An increase of their blood sugar levels (glucose level > 175 mg/dl) was considered diabetic and was used for the study.

### **Administration of the Test drug and the Control**

The test animals were divided into four treatments with 4 replicates. Treatment 1 (treated with 14.1 ml/kg of serpentina leaf extract), Treatment 2 (treated with 28.2 ml/kg of serpentine leaf extract), Treatment 3 (treated with 47 ml/kg of serpentine leaf extract) and Treatment 4 (treated with metformin as the positive control). The leaf extract was administered orally once and effects on blood glucose levels were measured 3 hours and 24 hours post medication.

### **Evaluation of biochemical parameter**

Blood glucose was measured with Accu-check glucometer using the tail prick method. Samples were taken before an initial 3 hours fasting and 6 days of alloxan administration. Blood glucose levels > 175 were considered hyperglycemic. Test animals with hyperglycemia were treated with the test extract and the control, and their glucose levels were measured 3 hours and 24 hours thereafter.

### **Statistical analysis**

The paired t-test was used to compare the anti-hyperglycemic effect of the test extract before and after test at 5% level of significance.

## RESULTS AND DISCUSSION

Analysis of the blood glucose profile of mice medicated with Alloxan showed a significant difference before and after administration of the drug.

### **Initial and Final Blood Glucose Levels of Mice after Medication with Alloxan**

The mean difference of the blood glucose levels after 3 and 24 hours post treatment were determined. Blood glucose levels were reduced with administration of metformin at a rate of 447 ml/kg with a mean difference of 18.75 after three (3) hours and a mean difference of ninety-three after twenty-four (24) hours. Administration with Serpentina leaf extract reveals a mean difference of reduction to -44.75, 69.45 and 140.5. The reduction was further observed after 24 hours of treatment with a mean difference obtained at 193.75, 233.75 and 264.5, respectively.

### **Mean Difference of Blood Glucose Levels after Treatments**

Hyperglycemia is the hallmark of diabetes and as a common chronic disease, it is reported that diabetes mellitus is becoming the third “killer” of humans along with cancer, cardiovascular and cerebrovascular diseases. Hyperglycemia happens when there is no insulin in the blood, not enough insulin in the blood, or the insulin in the blood is not working properly. The condition needs to be treated immediately. Once diagnosed, it is usually regulated by means of various commercially available drugs for its treatment.

However, prominent side-effects of such drugs are the main reason for an increasing number of people seeking alternative therapies that may have less severe or no side effects. Tiredness, dizziness, chills, blue/cold skin, muscle pain, difficulty in breathing, irregular heartbeat, stomach pain with nausea, vomiting or diarrhea, kidney complications, metal taste, skin rash, itching, weight gain, risk of liver disease, anemia, swelling of legs or ankles and hypoglycemia are the most serious known side effects.

In this study, the researchers investigated the anti-hyperglycemic activity of aqueous extract of *Serpentina* leaves which have been used by locals for the traditional management of diabetes with claims of efficacy. Our data showed that the reduction of blood glucose level of mice after three (3) and twenty-four (24) hours post treatment revealed a significant difference at a dose of 28.2 ml/kg *Serpentina* leaf extract. The foregoing findings only corroborates reports about the medicinal properties of *Serpentina* whose intake have shown favorable effects to people with diabetes.

The above-mentioned effect of *Serpentina* plant may be attributed to andrographolide that has been reported to have a wide range of biological activities, such as those that are anti-inflammatory, anti-allergic, anti-platelet aggregation, hepatoprotective, and anti- HIV. Andrographolide enhances the surface uptake of glucose by adipose tissues and inhibit glucose absorption from intestine and glucose production from liver. Review of literature also shows that some flavonoids and saponins isolated from medicinal plants significantly reduce blood sugar levels. Flavonoids stimulate the secretion of insulin in B-cells of islet of Langerhans of pancreas. It is possible that the presence of glycosides, flavonoids and tannins are responsible for their activity (Thanasekaran et. al, 2012).

### **Implications on Science Education**

The study will not only be useful for people with diabetes who cannot afford expensive treatments, but results can be used as data for the discussion of the health uses of *Serpentina* in medical and biology classes.

### **CONCLUSIONS**

Based on the results, the researchers conclude that a dose of 28.2 ml/kg *Serpentina* leaf extract showed favorable reduction of hyperglycemia in mice sustained after three (3) and twenty four (24) hours post treatment. Data further indicate that *Serpentina* (*Andrographis paniculata*) exhibited anti-hyperglycemic activity, with

effective glucose tolerance improvement capability compared to Metformin.

### RECOMMENDATIONS

Based on the results of the study, the researchers recommend the following:

1. A parallel study be conducted using other parts of the *Serpentina* (*Andrographis paniculata*) plant (i.e. Roots, barks, fruits and flowers)
2. Toxicity studies of *Serpentina* leaf extract should be conducted to determine the safety of the test drug.

### References

- Anju, D., Jugnu, G., Kavita, S., Arun, N., & Sandeep, D. (2012). A review on medicinal prospectives of *Andrographis paniculata* Nees. *Journal of Pharmaceutical and Scientific Innovation (JPSI)*, 1(1). [http://jpsionline.com/admin/php/uploads/3\\_pdf.pdf](http://jpsionline.com/admin/php/uploads/3_pdf.pdf)
- Chauhan, A., Sharma, P. K., Srivastava, P., Kumar, N., & Dudhe, R. (2010). Plants having potential antidiabetic activity: A review. *Der Pharmacia Lettre*, 2(3), 369-387. <https://www.scholarsresearchlibrary.com/articles/plants-having-potential-antidiabetic-activity-a-review.pdf>
- Gray, A. M., & Flatt, P. R. (1998). Antihyperglycemic actions of *Eucalyptus globulus* (*Eucalyptus*) are associated with pancreatic and extra-pancreatic effects in mice. *The Journal of Nutrition*, 128(12), 2319-2323. <https://doi.org/10.1093/jn/128.12.2319>
- Kumar, A., Dora, J., Singh, A., & Tripathi, R. (2012). A review on king of bitter (*Kalmegh*). *International Journal of Research in Pharmacy and Chemistry*, 2(1), 116-24. <https://doc.presentica.com/11870797/5ec20e2dd550f.pdf>

Rohilla, A., & Ali, S. (2012). Alloxan induced diabetes: Mechanisms and effects. *International Journal of Research in Pharmaceutical and Biomedical Sciences*, 3(2), 819-823. <https://bit.ly/3Ty9dyK>